

This is a repository copy of *Mirror and triplet displacement energies within nuclear DFT: : numerical stability*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/111953/>

Version: Accepted Version

---

**Article:**

Baczyk, Pawel, Dobaczewski, Jacek Jan [orcid.org/0000-0002-4158-3770](https://orcid.org/0000-0002-4158-3770), Konieczka, Maciej et al. (3 more authors) (2017) Mirror and triplet displacement energies within nuclear DFT: : numerical stability. *Acta Physica Polonica B*. pp. 259-264. ISSN 0587-4254

<https://doi.org/10.5506/APhysPolB.48.259>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Mirror and triplet displacement energies within nuclear DFT: numerical stability\*

P. BĄCZYK<sup>1</sup>, J. DOBACZEWSKI<sup>1,2,3,4</sup>, M. KONIECZKA<sup>1</sup>,  
T. NAKATSUKASA<sup>5,6</sup>, K. SATO<sup>7</sup>, W. SATUŁA<sup>1,4</sup>

<sup>1</sup>Faculty of Physics, University of Warsaw, Pasteura 5, PL-02-093 Warsaw, Poland

<sup>2</sup>Department of Physics, University of York, Heslington, York YO10 5DD, UK

<sup>3</sup>Department of Physics, P.O. Box 35 (YFL), FI-40014 University of Jyväskylä,  
Finland

<sup>4</sup>Helsinki Institute of Physics, P.O. Box 64, FI-00014 University of Helsinki,  
Finland

<sup>5</sup>Center for Computational Sciences, University of Tsukuba, Tsukuba 305-8577,  
Japan

<sup>6</sup>RIKEN Nishina Center, Wako 351-0198, Japan

<sup>7</sup>Department of Physics, Osaka City University, Osaka 558-8585, Japan

Isospin-symmetry-violating class II and III contact terms are introduced into the Skyrme energy density functional to account for charge dependence of the strong nuclear interaction. The two new coupling constants are adjusted to available experimental data on triplet and mirror displacement energies, respectively. We present preliminary results of the fit, focusing on its numerical stability with respect to the basis size.

PACS numbers: 21.10.Hw, 21.60.Jz, 21.30.Fe, 21.10.Dr

## 1. Introduction

An accurate description of atomic nucleus, a system of protons and neutrons interacting with electromagnetic and strong forces, is a difficult task. It can be simplified considerably by introducing the concept of isospin symmetry [1] that relies on charge independence, that is, on equality of nucleon-nucleon (NN) forces  $V_{pp} = V_{pn} = V_{nn}$  in the same space-spin channel. The NN scattering experiments indicate, however, that the strong interaction depends slightly on a pair of nucleons involved in the process [2]. On a fundamental level, violation of the isospin symmetry is due to the mass

---

\* Presented at the Zakopane Conference on Nuclear Physics Extremes of the Nuclear Landscape, Zakopane, Poland, August 28 – September 4, 2016

splitting and different charges of the *up* and *down* quarks and the difference in quark composition of proton and neutron.

In atomic nuclei, the main source of isospin-symmetry breaking (ISB) is the Coulomb force, which shifts binding energies of nuclei forming a multiplet of a given isospin  $T$ . This property is used to construct various mass indicators which are sensitive to the ISB effects. The most common among such indicators are the mirror (MDE) and triplet (TDE) displacement energies involving data on isospin doublets ( $T = \frac{1}{2}$ ) and triplets ( $T = 1$ ). It turns out, however, that the Coulomb interaction alone is not sufficient to fully explain neither the MDEs nor the TDEs and the additional ISB mechanism due to the strong nuclear force is needed [3, 4].

Contemporary *ab initio* models are able to account for ISB effects in both NN scattering data and light nuclei [5, 6, 7]. However, they are still not suitable for describing heavier systems which is a domain of mean-field (MF) or density functional theory (DFT). These approaches are excellent tools to study bulk properties (masses, radii or quadrupole moments) in atomic nuclei regardless of their mass and parity of proton and neutron numbers, see [9] and refs. cited therein. Among different variants of MF or DFT approaches the models based on the Skyrme interaction [8] are the most efficient computationally and fairly well describe nuclear binding energies. However, the isospin invariant Skyrme energy density functionals (S-EDF), which are typically used in practical applications, systematically fail to reproduce the experimental data on MDEs and TDEs. In our recent work [10] we introduced two new ISB terms into the S-EDF. The first calculations performed with these modifications proved the ability of the extended model to correctly grasp the missing ISB effects in both the MDEs and TDEs, however, a systematic fitting of the new coupling constants  $t_0^{\text{II}}$  and  $t_0^{\text{III}}$  is necessary.

In this paper we present a discussion on the numerical stability of MDEs and TDEs with respect to the choice of the basis size. The study allows us to estimate a theoretical uncertainty associated with a given basis cut-off. The discussion is followed by a presentation of preliminary results of the fitting procedure performed with a certain basis cut-off.

## 2. Numerical stability

Nuclear calculations often depend on a choice of the basis size. Their reliability and, in particular, predictive power requires an estimate of theoretical uncertainties related to the basis-size. The HFODD code [11] used in this work solves the HF equation in the Cartesian harmonic oscillator (HO) basis. Its size can be controlled by providing a number of spherical HO shells  $N$ . In practical applications, the choice of  $N$  is always a matter of trade-off

between computation time and expected precision of the calculations.

To evaluate theoretical uncertainty of the calculated MDEs and TDEs due to the basis size we have performed test calculations for  $T = \frac{1}{2}$  doublets with  $A=25, 33, 37, 57$ , and  $75$  and for  $T = 1$  triplets with  $A = 22, 34$ , and  $58$ . Calculations have been performed using the spherical HO bases consisting of  $N = 10, 12, 14$ , and  $16$  shells. For the heaviest doublets and triplets with  $A = 57, 58$ , and  $75$ , we have extended the test by including the bases consisting of  $N=18$  and  $20$  shells. In each case we have computed both the MDEs and TDEs using three different S-EDF parametrizations, including the density-independent  $SV_T$  S-EDF of Refs. [12, 13] and two popular density-dependent S-EDFs SkM\* [14] and SLy4 [15]. Results are collected in Figs. 1 and 2.

The basis-size-dependence tests suggest that the optimal strategy regarding both efficiency and accuracy of the calculations is to compute light ( $A = 10 - 30?$ ), medium-mass nuclei ( $A = 31? - 56$ ), and heavy ( $A \geq 57$ ) nuclei using bases consisting of  $N=10, 12$ , and  $14$  spherical HO shells, respectively. Both for the MDEs and TDEs, this strategy would result in the basis-size related uncertainty not exceeding  $\Delta_{\text{basis}} \approx \pm 15$  keV. As we show in Sect. 3, the basis-size uncertainty is relatively small as compared to the uncertainty resulting from the fitting procedure, which may justify using even smaller bases.

### 3. Fitting results

As stated in Sec. 2, the optimum choice of the number of spherical HO shells implies dividing the mass region of interest into the three subsets. In the present preliminary study, we use  $N=10$  shells for all nuclei having masses  $A = 10 - 56$  and  $58$ , and we use  $N = 14$  shells only for the heaviest systems. This choice almost doubles the basis-size related uncertainty of the calculated MDEs to  $\Delta_{\text{basis}} \approx \pm 30$  keV without much affecting the uncertainty of the calculated TDEs. A smaller computational time allows us to explore the richness of experimental data available for ISB effects in the  $A = 10 - 58$  nuclei. In future, it may allow us to perform calculations of mirror (MED) and triplet (TED) energy differences in rotational bands or the ISB effects in electromagnetic and  $\beta$  decays using the recently developed no-core configuration-interaction (NCCI) formalism [16], which involves CPU time demanding isospin and angular momentum projections and configuration mixing.

With the less strict choice of the basis size, the two new coupling constants (see Ref. [10]) were adjusted to all available data on MDEs ( $A = 10 - 75$ ) and TDEs ( $A = 10 - 58$ ). The fitting procedure has been realized independently for  $t_0^{\text{II}}$  and  $t_0^{\text{III}}$  parameters using linear regression method




Fig. 1. (Color online) Values of MDEs in function of the number of spherical HO shells used in the basis, plotted with respect to that obtained for  $N = 12$ . Results for multiplets with different values of  $A$  are labelled with different symbols, as shown in the legend. Panels (a), (b), and (c) show values obtained using  $SV_T$ ,  $SkM^*$ , and  $SLy4$  S-EDFs, respectively.

and following the guidelines from Ref. [17] and will be described in details in our forthcoming publication. The fitting results as well as the standard deviations from the experimental data points are presented in Table 1.

As it turns out, the uncertainty resulting from the choice of the basis cut-off,  $\Delta_{\text{basis}}$ , is small with respect to the values of standard deviations,  $\sigma_{\text{fit}}$ , given in Table 1. Hence, in the total uncertainty,  $\Delta_T = \sqrt{\sigma_{\text{fit}}^2 + \Delta_{\text{basis}}^2}$ , the basis-size related uncertainty constitutes only a small correction.




Fig. 2. (Color online) Same as in Fig. 1, but for the TDEs.

Table 1. Values of the  $t_0^{\text{II}}$  and  $t_0^{\text{III}}$  parameters with uncertainty and standard deviations for MDE and TDE,  $\sigma_{\text{fit}}$ , resulting from the fit to all available data on isospin doublets and triplets. The calculations has been done for three different S-EDFs considered in this work.

Interaction	SV	SkM*	SLy4
$t_0^{\text{II}}$ (MeV fm <sup>3</sup> )	$17 \pm 5$	$24 \pm 8$	$22 \pm 7$
$\sigma_{\text{fit}}$ (keV)	100	110	100
$t_0^{\text{III}}$ (MeV fm <sup>3</sup> )	$-7.3 \pm 1.9$	$-5.5 \pm 1.3$	$-5.5 \pm 1.1$
$\sigma_{\text{fit}}$ (keV)	190	150	120

#### 4. Summary

In the paper we performed preliminary calculations of MDEs and TDEs paying special attention to the numerical stability of the results with respect

to the basis size. It turned out that the optimum choice of the number of spherical HO shells,  $N$ , which defines the basis size, is  $N = 10$  for light nuclei ( $A = 10 - 30$ ?),  $N = 12$  for medium-mass nuclei ( $A = 31 - 56$ ) and  $N = 14$  for heavy nuclei ( $A \geq 57$ ). It is shown, that the corresponding uncertainty,  $\Delta_{\text{basis}} \approx 15 \text{ keV}$  is small as compared to the uncertainty coming from the fitting procedure. This allows us to compute light and medium-mass nuclei using the basis consisting only of  $N = 10$  HO shells without losing much of the precision. Smaller basis, in turn, will be beneficial regarding efficiency of the planned NCCI calculations. Finally, the two new coupling constants were adjusted to all available experimental data. The details of the fitting procedure will be presented in the forthcoming publications.

This work has been supported by the Polish National Science Centre under the contract 2015/17/N/ST2/04025 and by the Academy of Finland and University of Jyväskylä within the FIDIPRO program. Advanced computational resources were provided by the Świerk Computing Centre (CIS) at the National Centre for Nuclear Research (NCBJ).

## REFERENCES

- [1] E. Wigner, Phys. Rev. **51**, 106 (1937).
- [2] G.A. Miller, B.M.K. Nefkens, and I. Šlaus, Phys. Rep. **194**, 1 (1990).
- [3] J.A. Nolen and J.P. Schiffer, Ann. Rev. Nuc. Sci. **19**, 471 (1969).
- [4] W. Satuła, J. Dobaczewski, M. Konieczka, and W. Nazarewicz, Acta Phys. Polon. B **45**, 167 (2014).
- [5] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
- [6] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
- [7] R.B. Wiringa *et al.*, Phys. Rev. C **88**, 044333 (2013).
- [8] T.H.R. Skyrme, Nucl. Phys. **9**, 615 (1959).
- [9] M. Bender, P.-H. Heenen, and P.-G. Reinhard, Rev. Mod. Phys. **75**, 121 (2003).
- [10] P. Bączyk, J. Dobaczewski, M. Konieczka, and W. Satuła, Acta Phys. Pol. B Proc. Suppl. **8**, 539 (2015).
- [11] N. Schunck *et al.*, Comput. Phys. Commun. **183**, 166 (2012).
- [12] M. Beiner, H. Flocard, N. Van Giai, and P. Quentin, Nucl. Phys. A **238**, 29 (1975).
- [13] W. Satuła, J. Dobaczewski, W. Nazarewicz, and M. Rafalski, Phys. Rev. Lett. **106**, 132502 (2011).
- [14] J. Bartel, P. Quentin, M. Brack, C. Guet, and H.B. Håkansson, Nucl. Phys. A **386**, 79 (1982).
- [15] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer, Nucl. Phys. A **635**, 231 (1998).

- [16] W. Satuła, P. Bączyk, J. Dobaczewski, and M. Konieczka, Phys. Rev. C **94**, 024306 (2016).
- [17] J. Dobaczewski, W. Nazarewicz, and P.G. Reinhard, J. Phys. G: Nucl. and Part. Phys. **41**, 074001 (2014).